

CITY OF YUCAIPA
ORDINANCE NO 347
POULTRY RANCH INSPECTION

Ranch Name: HOOVER EGG RANCH – 34618 AVE H

Date: September 1, 2017

PROPERTY MAINTENANCE

WATER SYSTEM/COOLING SYSTEM:	YES	NO
Leaks present	X	
Leaks required to be repaired	N/A <input type="checkbox"/> IMMEDIATELY <input checked="" type="checkbox"/>	24 HOURS <input type="checkbox"/>
Leaks causing wet manure	X	
Wet manure being removed or treated with lime	X	
FEED:	YES	NO
Spilled feed present		X
Evidence of rodent Infestation present		X
EQUIPMENT:	YES	NO
Operable spray equipment on site	X	

MANURE DISPOSAL, DEAD BIRDS, BROKEN EGGS

MANURE:	YES	NO
Raw Manure removed from the ranch within three (3) days	X	
DEAD BIRDS/BROKEN EGGS:	YES	NO
Stored in Fly-tight containers	X	
Removed from ranch weekly	X	

FLY CONTROL

BAIT STATIONS:		YES	NO	
Adult fly bait station located at the end of every house		X		
Bait stations maintained properly		X		
Adult flies:	NOT OBSERVED	LIGHT	MODERATE	HEAVY
Adult fly population on the ranch			X	
Emergency Activities need to be implemented		YES	NO	X

LARVA BREEDING:	NOT OBSERVED	LIGHT	MODERATE	HEAVY
Larva breeding on the ranch	X			
Emergency activities need to be implemented		YES	NO	X

RANCH IS IN COMPLIANCE	YES	X	NO
-------------------------------	-----	---	----

OBSERVATIONS:

Fly population is about the same as last inspection. Majority of dry manure has been removed from under the coops. No larvae noticed during inspection of several different locations. Owner sprayed twice this week (Tuesday and Thursday). Several inch thick layer of manure observed where piles were placed for pick up. Evidently new driver for manure removal didn't scrape down far enough to remove all manure residue. Ranch owner will make sure that is taken care of during next pick up (Saturday). Several chicken parts and broken eggs were also noted but were less than last inspection and were removed during inspection.

RECOMMENDATIONS:

Continue to remove chicken manure per the Manure Management Plan requirements. Place soil over remaining dry manure to encourage biological insects that aid in abating larvae and to bring grade up for easy removal of fresh manure. Continue with the spraying 2 times a week. Continue to abate areas between coops of feathers, eggs, and vegetation (especially vegetation-should be no higher than 4 inches).

ACTION TAKEN:

Install at least 2 inches of dirt over the dry (non-larvae breeding) manure. Fix all leaks in cups and mister system (ongoing issue).

Biosecurity Measures Taken Yes x No

Inspected By: P. Swantek